

Radiosonde data denial studies and implications for height attainment

GRUAN ICM-15, 14 March 2024

Bruce Ingleby and Inna Polichtchouk

ECMWF

Bruce.Ingleby@ecmwf.int

With thanks to Sean Healy and Tony McNally



Overview

- Seasonality of stratospheric variability and FSOI (recap from Nov 2021)
- Results from data denial studies (OSEs)
 - Control (all data)
 - noRS – no radiosonde data
 - noRSlt100 – no radiosonde data for $P < 100$ hPa
 - noRStq – no radiosonde temperature or humidity
- Geostrophic balance in the stratosphere
- Other: Modern O-B statistics, assimilation of stratospheric humidity?
- Summary

Stratospheric seasonality


- work with Inna Polichtchouk
U10 hPa @ 60°N, 60°S

- Different years plotted 1950 to 2021
- Data from ERA5 (Hersbach et al, QJ, 2020)
- Winter: strong polar vortex
- SH vortex stable except when breaking down in Austral spring
- In NH winter planetary waves disturb the polar vortex
- Largest disturbances form stratospheric sudden warmings (SSWs)
- In summer there are about 4 months when nothing much happens – except a few gravity waves




Annual cycle of radiosonde impact in stratosphere 45-90N

- Forecast Sensitivity to Observation Impact (FSOI) – estimate of the importance of data in reducing errors in T+24 h forecasts (doesn't account for anchoring in bias correction)
- Radiosonde FSOI for different variables: 4 years
- Wind (U,V) FSOI > temperature FSOI
- Larger values Nov-April despite lower observation numbers (green bars)
 - Balloons burst at lower altitudes in winter ☹️
 - Especially DJF
 - Could use larger balloons then (cost)
- Somewhat noisy plots ...
- Upper level data in NDJF 'worth' about 3* the data in MJJA by this measure! See also OSEs



Annual cycle in (median) burst height

- Seasonality in burst height is largest at high latitudes
- Can see effect of larger balloon at Sodankyla (blue dashed line) – selected months
- Lower plot:
 - Alaska: 600 g balloons
 - Canada: 800 or 600 g
 - Russia: 500 g
 - Europe: varies, mainly 350 and 600 g
- Gas used will also affect height




T+12 fit to radio occultation (RO) data when removing sondes

- In NH ~60% larger impact in DJF; minor impact of temperature (red line)


Relative RMS(O-B) fit to RO for radiosonde OSEs 20-90N

Relative RMS(O-B) fit to RO for radiosonde OSEs 20S-20N





Wind impact from T+12 to day 9: noRS and noRSIt100

- Vs operational analyses
- Largest impact at short-range in NH (expected)
- Largest impact 150-50 hPa
- noRSIt100: impact mainly local, stays at P<100 hPa
- Larger impact of RS winds than temperatures partly because there are plenty of satellite data giving mass information
- Few satellite winds, esp. at upper levels



Timeseries vs forecast range


- ~20% impact at 100 and 50hPa in NH from noRS
- At 50 hPa and higher the **It100** results match the noRS (black) results (right-hand panel)






Geostrophic balance at 20 hPa, op T+24 fcs on 1x1 deg grid

- Winter mid-latitudes close to geostrophic balance
- Also very strong winds
- The balance means that satellite data help with the winds more in winter

Dec 2022 - Jan 2023 T+24 20 hPa





June/July 2023 T+24 20 hPa




Misc 1. Modern O-B statistics vs RS41, standard levels, 2023

- 20-90N, T and wind (not shown) OK; RH, M20 has bias up to 7.5% at 200 hPa
- RS41 presumed ~unbiased, known that model is too wet just above tropopause



Misc 2: RS41 RH Summer 2023 by latitude band: 150-10 hPa

- To date ECMWF has assimilated humidity in the troposphere but not the stratosphere
- Starting to plan for assimilation of lower stratospheric humidity
 - Help with model biases
 - Hunga-Tonga eruption!
- Hence look at RS41 humidity:
- Mostly usable (if sparse)? 😊
- Check model T-q error correlation
- Main data problem appears to be in profiles from Antarctica – would omit those



Stratospheric seasonality summary

- Much more variability in NH winter (SH spring) stratosphere than in summer
- Radiosondes have much larger FSOI <100 hPa in Winter, mainly from winds
- OSEs show smaller seasonality, but more impact in Winter, mainly from winds
- Satellite temperature data very important – helps more with the winds in Winter: approximate geostrophic balance then
- Removing sonde data at $P < 100$ hPa impact is largely local
- Larger RS impact in Winter is despite lower burst height then – case for larger balloons + more gas in Winter